

The spectral type of sums of operators on non-Hilbertian Banach lattices

Ian Doust and Gilles Lancien

Abstract

It is known that on a Hilbert space the sum of a well-bounded operator and a commuting real scalar-type spectral operator is well-bounded. It had been conjectured that this may still hold for operators on L^p spaces for $p \neq 2$. We show here that this conjecture is false. ¹

1 Introduction

There is by now a significant body of results which relate the spectral decomposition properties, or the functional calculus properties, of the sum of two commuting operators to those of the summands. Two much-studied classes are the scalar-type spectral operators and the well-bounded operators. For simplicity, let X be a reflexive Banach space. On such a space, an operator T is scalar-type spectral if and only if it admits an integral representation with respect to a countably additive projection-valued measure, or equivalently, if it admits a $C(\sigma(T))$ functional calculus. An operator $T \in B(X)$ is well-bounded if it admits a Riemann-Stieltjes type integral representation with respect to an increasing family of projections on X , or, equivalently, if it admits an $AC[a, b]$ functional calculus. (Fuller definitions are given in

¹2000 Mathematics Subject Classification 47B40 (primary)

Section 2.) The distinction between these classes is that the spectral expansions for a scalar-type spectral operator are of an unconditional nature, while those for well-bounded operators correspond to conditional decompositions of the Banach space.

C. A. McCarthy [11, 12] proved that the sum and product of two commuting scalar-type spectral operators acting on an L^p ($1 \leq p \leq \infty$) space is again scalar-type spectral. This result has been extended to a wide class of other Banach spaces, including, for example, weakly complete Banach lattices (see [5]). On the other hand, there are reflexive spaces (indeed even UMD spaces), such as the von Neumann-Schatten C_p spaces, on which the sum of two commuting scalar-type spectral operators may fail to be scalar-type spectral. More recently, it was shown in [2] that on spaces with Kalton and Weis' property (Δ), and in particular on all UMD spaces, the sum of two commuting scalar-type spectral operators was however always well-bounded. As was noted in [2], this result can not be extended to all reflexive spaces.

T. A. Gillespie [4] showed that even on ℓ^2 , the sum of two commuting well-bounded operators need not be well-bounded. In a positive direction however, he showed that on this space, the sum and product of a real scalar-type spectral operator (that is, a scalar-type spectral operator with real spectrum) and a commuting well-bounded operator are always well-bounded. It had been asked whether this last result might hold on a wider class of spaces such as the reflexive L^p spaces (see, for example, [2, Question 2.7]). In [2] it was shown that a sufficient condition for the sum of a real scalar-type spectral operator S and a commuting well-bounded operator T to be again well-bounded is that the spectral family of projections associated with T be R -bounded.

The aim of this paper is to note that part of the construction given by N. J. Kalton and the second author in [6] can be easily adapted to produce counterexamples to the above question on any non-Hilbertian reflexive

Banach lattice.

2 Well-bounded operators

An operator $T \in B(X)$ is well-bounded if it admits an $AC(J)$ functional calculus for some compact interval $J \subset \mathbb{R}$. Here $AC(J)$ denotes the absolutely continuous functions on $J = [a, b]$, which is a Banach algebra under the norm $\|f\|_{AC(J)} = |f(b)| + \text{var}_J f$. Thus, T is well-bounded if there exists some constant K and a compact interval J such that $\|f(T)\| \leq K \|f\|_{AC(J)}$. We shall use the notation

$$\|T\|_{WB} := \sup\{\|f(T)\| : f \in AC[a, b] \text{ and } \|f\|_{AC(J)} = 1\}.$$

Standard results about well-bounded operators ensure that this quantity depends only on T and not on the choice of interval.

Associated to each well-bounded operator on a reflexive Banach space is a uniformly bounded family of projections known as a spectral family. If a well-bounded operator T has associated spectral family $\{E(\lambda)\}_{\lambda \in \mathbb{R}}$, then for all $f \in AC(J)$,

$$f(T) = \int_J^\oplus f(\lambda) dE(\lambda).$$

The integral here is a Riemann-Stieltjes type integral.

We refer the reader to [3] for further details (and the appropriate definitions for nonreflexive spaces) of the theory of well-bounded and scalar-type spectral operators. Background definitions concerning bases in Banach spaces and Banach lattices may be found in [8] and [9].

3 The main result

The first and main step is to consider the case of a reflexive Banach space with an unconditional basis.

Theorem 3.1. *Let X be a reflexive Banach space which is not isomorphic to ℓ^2 . If X admits an unconditional basis, then there exist a real scalar-type spectral operator $S \in B(X)$ and a well-bounded operator $T \in B(X)$ such that $ST = TS$ and $S + T$ is not well-bounded.*

Proof. It was shown in [10] (see the remark at the end of that paper) that if a Banach space X has a normalized unconditional basis and every normalized unconditional basis of X is symmetric, then X is isomorphic to either c_0 , ℓ^1 , or ℓ^2 . It follows that if a reflexive space X which is not isomorphic to ℓ^2 has an unconditional basis, then X admits a normalized unconditional basis which is not symmetric. In particular X must admit a normalized unconditional basis (x_n) which is not equivalent to any of the standard bases of c_0 or ℓ^p ($1 \leq p < \infty$).

The proof of Theorem 4 of [7] (see also Theorem 2.a.10 of [8]) then shows that there exists a permutation π and a block basis (v_n) of $(x_{\pi(n)})$ whose closed linear span is not complemented. To simplify the notation we may assume that (v_n) is in fact a block basis of (x_n) , with

$$v_n \in \text{span}(x_{r_n+1}, \dots, x_{r_{n+1}}),$$

where (r_n) is a suitably chosen increasing sequence of integers. Let $X_n = \text{span}(x_{r_n+1}, \dots, x_{r_{n+1}})$, and let P_n denote the projection from X onto X_n determined by the basis projections. Let $\alpha_n = \frac{1}{n^2}$. The unconditionality of (x_n) implies that $S := \sum_{n=1}^{\infty} \alpha_n P_n$ is a real scalar-type spectral operator on X .

For each n , there is a norm one projection \tilde{V}_n from X_n onto the span of v_n . Let $V_n = \tilde{V}_n P_n$ and $W_n = P_n - V_n$. Note that $(V_1, W_1, \dots, V_n, W_n, \dots)$ defines a Schauder decomposition of X .

Let $Q_1 = V_1$ and for $n \geq 2$, let $Q_n = W_{n-1} + V_n$. Then $\{Q_n\}$ is a sequence of disjoint finite-rank projections. Note that for all N ,

$$\sum_{n=1}^N Q_n = V_N + \sum_{n=1}^{N-1} P_n$$

and so these partial sums are uniformly bounded. Let $\beta_n = -\frac{1}{n^2} - \frac{1}{n^3}$. It follows from [1, Theorem 3.3] that $T := \sum_{n=1}^{\infty} \beta_n Q_n$ converges to a well-bounded operator on X .

It is straightforward to verify that for all m and n , P_m and Q_n commute. It follows that S and T commute. It remains then to show that $S + T$ is not well-bounded. Suppose, to the contrary, that $S + T$ is well-bounded.

A small calculation shows that

$$S + T = \sum_{n=1}^{\infty} -\frac{1}{n^3} V_n + \sum_{n=1}^{\infty} \gamma_n W_n,$$

where $\gamma_n = \alpha_n + \beta_{n+1}$ is positive for all $n \geq 1$. Consider now, for $n \geq 1$, $f_n \in AC[-1, 1]$ chosen so that $f_n \equiv 1$ on $[-1, -\frac{1}{n^3}]$, $f_n \equiv 0$ on $[-\frac{1}{(n+1)^3}, 1]$ and f_n decreases linearly on $[-\frac{1}{n^3}, -\frac{1}{(n+1)^3}]$, then $f_n(S + T) = \sum_{j=1}^n V_j$. We now claim that the increasing sequence of projections $(\sum_{j=1}^n V_j)_{n=1}^{\infty}$ is unbounded. Otherwise, since X is reflexive, it would converge in the strong operator topology and its limit would be a bounded projection onto the closed linear span of $\{v_n, n \geq 1\}$. Thus $\{\|f_n(S + T)\|\}$ is unbounded and, as $\|f_n\|_{AC(J)} = 2$ for each n , it follows that $S + T$ must not be a well-bounded operator.

□

The hypotheses in this theorem are not optimal, but they do suffice to cover the cases of most interest. In particular, we have as a first corollary:

Corollary 3.2. *Let $1 < p < \infty$, $p \neq 2$, and let $X = L^p[0, 1]$. Then there exist a real scalar-type spectral operator $S \in B(X)$ and a well-bounded operator $T \in B(X)$ such that $ST = TS$ and $S + T$ is not well-bounded.*

Proof. The Haar system forms an unconditional basis for X . □

Finally we prove the result announced in the introduction.

Corollary 3.3. *Let X be a reflexive Banach lattice, which is not isomorphic to a Hilbert space. Then there exist a real scalar-type spectral operator $S \in B(X)$ and a well-bounded operator $T \in B(X)$ such that $ST = TS$ and $S + T$ is not well-bounded.*

Proof. We will essentially follow the proofs of Theorems 3.5 and 3.7 in [6]. Let us assume that for any real scalar-type spectral operator $S \in B(X)$ and any well-bounded operator $T \in B(X)$ commuting with S , $S + T$ is well-bounded. We will then show that X is order isomorphic to a Hilbert space. For this, since X is reflexive and therefore order continuous, it is enough to prove that every normalized sequence of disjoint elements of X is equivalent to the canonical basis of ℓ^2 (see [9] Lemma 1.b.13).

So, let (v_n) be a normalized sequence of disjoint elements of X . As in [6], we can pick an unconditional Schauder decomposition (X_n) of X such that for each n , X_n is an ideal of X and $v_n \in X_n$. Then, by repeating the argument in the proof of Theorem 3.1, we obtain that $Y = \overline{\text{span}}\{v_n, n \geq 1\}$ is complemented in X . Thus, for any real scalar-type spectral operator $S \in B(Y)$ and any well-bounded operator $T \in B(Y)$ commuting with S , $S + T$ is well-bounded (indeed, any counterexample on Y could easily be extended into a counterexample on X). Finally, it follows from Theorem 3.1 that (v_n) is equivalent to the canonical basis of ℓ^2 . \square

Remarks. (1) Virtually identical constructions provide examples where the product rather than the sum fails to be well-bounded; with $\{\alpha_n\}$, $\{\beta_n\}$, $\{P_n\}$, and $\{Q_n\}$ as in the proof of Theorem 3.1, then setting

$$S = I + \sum_{n=1}^{\infty} (e^{\alpha_n} - 1)P_n \quad \text{and} \quad T = I + \sum_{n=1}^{\infty} (e^{\beta_n} - 1)Q_n$$

will suffice.

(2) It should be noted that on nonreflexive spaces, these questions are somewhat more complicated since the relationship between the functional calculus conditions and the integral representation conditions is not as strong. On spaces such as ℓ^∞ , or indeed any Grothendieck space with the Dunford-Pettis property, the countable additivity requirement for a spectral measure allows very few scalar-type spectral operators. On such a space every scalar-type spectral operator is a finite linear combination of disjoint projections

[13] and consequently the sum of a real scalar-type spectral operator and a commuting well-bounded operator is always well-bounded. On other nonreflexive spaces it is relatively easy to construct explicit examples where the sum of a real scalar-type spectral operator and a commuting well-bounded operator is not well-bounded. For example, on the space of trace class operators C_1 , the operators $S(A) = D_1A$ and $T(A) = AD_2$, where D_1 and D_2 are diagonal matrices with real entries, are commuting scalar-type spectral operators whose sum, in general, will fail to be well-bounded. The situation on the classical sequence space c_0 and ℓ^1 appears, however, to be open.

(3) On nonreflexive spaces a more natural line of enquiry might be to seek conditions on Banach spaces which allow one to deduce functional calculus properties of the sum of commuting operators S and T from the functional calculus properties of the individual operators.

References

- [1] Cheng, Q. and Doust, I., Compact well-bounded operators, *Glasg. Math. J.* **43** (2001), 467–457.
- [2] Doust, I. and Gillespie, T. A., Well-boundedness of sums and products of operators, *J. London Math. Soc. (2)* **68** (2003), 183–192.
- [3] Dowson, H.R., Spectral theory of linear operators, London Mathematical Society Monographs 12, Academic Press, London, 1978.
- [4] Gillespie, T. A., Commuting well-bounded operators on Hilbert spaces, *Proc. Edinburgh Math. Soc. (2)* **20** (1976), 167–172.
- [5] Gillespie, T. A., Boundedness criteria for Boolean algebras of projections, *J. Funct. Anal.* **148** (1997), 70–85.

- [6] Kalton, N. J., Lancien, G., A solution to the problem of L^p -maximal regularity, *Math. Z.* **235** (2000), 559–568.
- [7] Lindenstrauss, J., and Tzafriri, L., On the complemented subspaces problem, *Israel J. Math.* **9** (1971), 263-269.
- [8] Lindenstrauss, J., and Tzafriri, L., Classical Banach Spaces I, Springer-Verlag, Berlin, 1977.
- [9] Lindenstrauss, J., and Tzafriri, L., Classical Banach Spaces II, Springer-Verlag, Berlin, 1979.
- [10] Lindenstrauss, J., and Zippin, M., Banach spaces with unique unconditional basis, *J. Funct. Anal.* **3** (1969), 115–125.
- [11] McCarthy, C. A., Commuting Boolean algebras of projections, *Pacific J. Math.* **11** (1961), 295–307.
- [12] McCarthy, C. A., Commuting Boolean algebras of projections. II. Boundedness in L_p , *Proc. Amer. Math. Soc.* **15** (1964), 781–787.
- [13] Ricker, W., Spectral operators of scalar type in Grothendieck spaces with the Dunford-Pettis property, *Bull. London Math. Soc.* **17** (1985), 268–270.

SCHOOL OF MATHEMATICS, UNIVERSITY OF NEW SOUTH WALES, SYDNEY, NSW 2052, AUSTRALIA

E-mail address: i.doust@unsw.edu.au

EQUIPE DE MATHÉMATIQUES UMR6623, UNIVERSITÉ DE FRANCHE-COMTÉ, F-25030 BESANÇON CEDEX, FRANCE

E-mail address: gilles.lancien@math.univ-fcomte.fr